



# TAKING THE MYSTERY OUT OF ORAL MEDICATIONS

BRAD SUTTON, OD, FAAO

CLINICAL PROFESSOR

INDIANA UNIVERSITY SCHOOL OF OPTOMETRY

# FINANCIAL DISCLOSURES

- No financial disclosures

## CATEGORIES

- I ) ANTIBIOTICS
- II ) ANTIVIRALS
- III ) PAIN RELIEF
- IV ) STEROIDS



## ORAL ANTIBIOTICS: OCULAR INDICATIONS

- Beat the bugs!
- Rosacea / Ocular Rosacea
- Dacryoadenitis
- Dacryocystitis
- Preseptal Cellulitis
- Hordeola / Chalazia
- Blowout Fractures



## COST

- \$4 (30 day) and \$10 (90 day) lists for generics at Kroger, Walmart, Target etc.
- Indicated by an \*
- Meijer has some generic antibiotics for free (Cipro, Cephalexin, Amoxicillin, PCN-VK, Ampicillin, SMZ-TMP)



## CEPHALEXIN \*

- 250 or 500 mg (QID or BID)
- Excellent broad spectrum cephalosporin
- Bactericidal
- Cross sensitivity with penicillin regarding allergies but not with everyone
- Keflex brand is very expensive!
- Up to 40% failure rate with facial cellulitis due to resistance

## DICLOXACILLIN

- Penicillinase resistant penicillin
- Great for soft tissue infections
- Bactericidal
- Nausea, allergies, diarrhea
- 250 mg QID or 500 mg BID
- Inexpensive

## AUGMENTIN

- Amoxicillin plus clavulanate: 250 ,500 mg TID or 875mg BID
- Works on bugs that are resistant to penicillin due to penicillinase
- Bactericidal, good coverage, allergies
- 500mg now available generically, but more expensive than dicloxacillin



## TETRACYCLINE\*

- 250 or 500 mg QID
- Bacteriostatic but much resistance
- Poor for soft tissue disease
- Can not be used in pregnant women or children due to effect on bone and enamel formation (discoloration of teeth)
- Makes BC Pill less effective. Yeast infections.
- Photosensitivity, stomach upset, calcium inactivation (take on empty stomach)
- Great lipid / acid modulating effects

## DOXYCYCLINE

- 50 or 100 mg BID
- Periostat: 20mg (mostly dental use)
- In tetracycline family
- Can take with food
- Less problems with photosensitivity
- Still get stomach upset
- As effective as tetracycline but fewer side effects, better dosing.
- Oracea (30 /10 ) \$\$\$\$\$\$\$\$\$\$\$\$ (very expensive)
- Doxy used to be inexpensive, but not now. Cost went up 10 fold. Used to be on \$4 plans, but no longer. Several generic makers stopped making it, cost went up
- Can also use minocycline

## AZITHROMYCIN

- Zithromax Z-pack: 6, 250 mg capsules. Is a macrolide.  
Moderate price
- Take 500 mg ( 2 ) the first day and one 250 mg tablet each of the next 4 days
- Can also take a single, 1000 mg dose (for chlamydia for example)
- May enhance the effect of oral anticoagulants
- Expensive but great for compliance
- 2 X risk of sudden cardiac death in heart patients

## AZITHROMYCIN

- Now has FDA warning for fatal arrhythmia
- Greater risk if prolonged QT interval, bradycardia, hypomagnesemia. Some studies refute this risk, so controversial
- Many experts calling for ban due to resistance concerns. Long half life and broad spectrum contribute majorly to overall resistance.
- Can be as effective in treating rosacea / MGD / chalazia as the tetracycline / doxycycline family of drugs

## ERYTHROMYCIN\*

- Ery-tab sustained release tablets 250, 333, or 500 mg. Dose is 1000 mg ( 1 gram) per day so dose according to tablet
- Can use safely when tetracycline family can not be used (children, etc.)
- Bacteriostatic and terrible stomach upset
- Dose not have the lipid / acid modulating properties of the tetracyclines
- Never a first choice

## ERYTHROMYCIN

- Increased risk of sudden cardiac death
- Two-fold increase of very low risk when taken alone
- Five-fold increase when taken with the following drugs.....  
Diltiazim, Fluconozole, Itraconozole, Ketaconozole, Verapamil
- These drugs slow the breakdown of E-mycin resulting in increased concentration which in turn increases cellular sodium levels in resting heart muscle cells triggering an arrhythmia



## BACTRIM

- Trimethoprim and Sulfamethoxazole: one tablet contains 80 mg T and 400 mg S (also available in double strength ). One double-strength tablet Q12h
- Can not use if patient has sulfa allergy
- Good against MRSA and toxoplasmosis

## CIPROFLOXACIN\*

- Fluoroquinolone: 750 / 500 / 250 BID
- 5mg/100ml suspension
- Effective but overused so resistance an issue. Lovaquin shows less resistance
- Can not use in patients under 18 due to joint / tendon problems
- Possible increased risk of RD? 3% vs. .6%. Other large studies refute this




## ORAL FLUOROQUINOLONES

- Significant side effects.....
  - Peripheral neuropathy
  - Tendon rupture
  - Heart arrhythmia
  - Dysglycemia in diabetics

## ORAL ANTIVIRALS

- Used to manage Herpes Simplex and Herpes Zoster



## ORAL ANTIVIRALS-DOSING SIMPLEX

- Acyclovir (200\*,400,800) : typically 800mg TID, varies
- Prophylactic dosing 400mg BID
- Also available in a pediatric suspension
- 200 mg available on \$4 / \$10 plans, but only allocated one 200 mg tablet per day, so problematic
- Famvir (125,250,500)
- 500mg TID
- Valtrex (500,1000)
- 500 mg TID



## ORAL ANTIVIRALS-DOSING ZOSTER

- Acyclovir: 800mg 5X day for 10 days
- Famvir: 500mg TID x 1week (may be antiviral of choice with zoster: can kill latent virus particles)
- Valtrex: 1000 mg TID X 1 week



## SIDE EFFECTS OF ANTIVIRALS

- Very safe
- Significant caution with renal impairment: only true contraindication other than allergy
- Headache
- GI upset / abdominal pain




## H.E.D.S. (HERPETIC EYE DISEASE STUDY) - FINDINGS

- Prophylactic 400 mg of oral Acyclovir (Famvir / Valtrex not studied) twice per day for one year resulted in a 45% decrease in the rate of recurrence for all forms of ocular complications
- Over the six months after discontinuation, there was no rebound increase but no continued benefit, so have to keep taking it
- Interestingly, the benefit mostly applied to those with previous stromal disease, not previous dendrites alone in this study

## RECENT STUDY

- Olmstead County, Minnesota (394 patients)
- Those NOT taking prophylactic antivirals were.....
- 9.4 X more likely to have epithelial recurrence
- 8.4 X more likely to have stromal rec.
- 34.5 X more likely to have lid / conj. rec



## PROPHYLAXIS

- So.....
- At least discuss prophylaxis for all patients with stromal disease and patients with multiple attacks of epithelial disease
- Acyclovir 400mg PO BID
- Very safe, caution in severe kidney disease



## PROPHYLAXIS

However: Report in Journal of Infectious Disease  
2013:208 (November) 1359-1365 and an editorial in the  
same issue.....

- Are we creating Acyclovir resistant strains of HSV with prophylactic use?
- In cases using Acyclovir for ocular prophylaxis, 26% showed ACV resistance. So now we must consider this

## ORAL PAIN MEDICATIONS

- Manage underlying condition appropriately first from an ocular standpoint
- Topical/ocular pain control.....
- Cycloplegia
- NSAIDS
- Steroids
- Bandage CL
- Topical anesthetic in office only

## PAIN MEDICATIONS

- If topical management is not enough, then consider oral pain relief
- Laws vary for OD's regarding use of controlled substances
- Two broad categories...
- OTC pain relief, mostly NSAID's
- Narcotic pain relief

## NSAIDS

- OTC NSAID's are often enough to mitigate ocular pain
- Aspirin, Ibuprofen, APAP, naproxen
- Common Trade names aspirin, Advil, Tylenol, Aleve
- Aspirin 81mg, 325-500mg
- Advil 200mg
- Tylenol 325-500mg
- Aleve 220mg

## RX NSAIDS

- Indomethacin (Indocin) 25, 50 mg
- Naproxen (Anaprox) 275, 550 mg
- Ibuprofen (Motrin) 200-800 mg
- Indomethacin very good for scleritis. TID dosing



## COMMON NSAID CONCERNS

- GI upset (take with food or drink, don't lie down for 30 minutes)
- Bleeding
- Ulcers
- Caution also with renal disease, heart disease, liver disease (mostly APAP)
- Rx strength particularly problematic with heart disease

## TRAMADOL (ULTRAM): USED TO BE NON-NARCOTIC, BUT NOW A CONTROLLED SUBSTANCE

- Immediate release (50-100 mg) and extended release (100-300 mg) versions
- Maximum dose 300mg /day
- Dose q 6-8 h
- Schedule IV, so limited (but possible) abuse potential





## NARCOTIC PAIN RELIEF

- As an OD, may or may not have authority to use (only Tramadol in Indiana)
- Standard warnings.....no alcohol, don't operate machinery





## NARCOTIC SIDE EFFECTS

- Constipation very common, and can be severe
- Nausea and vomiting: often ceases after first few doses
- Sedation
- Lack of mental clarity
- Respiratory depression (most severe)

## NARCOTIC PAIN RELIEF

- DEA Scheduled substances
- I-V
- Schedule one has high abuse potential, schedule 5 very limited abuse potential
- Two types of dependence....
- Psychological and physical
- Physical usually requires 2 weeks of therapy or more

## OXYCODONE

- Schedule II :high abuse potential with severe dependence risk
- Percocet: 5mg with 325 mg of APAP
- Percodan:4.5mg with 325 of APAP



- Tylox: 5mg with 500mg of APAP

## HYDROCODONE

- Schedule II now
- Lortab: 2.5,5,7.5 mg with 500mg APAP
- Vicodin: 5mg with 500mg APAP
- Vicodin ES: 7.5MG with 500mg APAP
- Norco: 5,7.5,10 with 325 APAP
- Zohydro ER: 10,15,20,30,40,50



## CODEINE

- Schedule III
- Tylenol with Codeine, all have 300mg of APAP
- Tylenol #2 : 15mg
- Tylenol #3 : 30mg
- Tylenol # 4 : 60 mg



## ORAL STEROIDS

- When oral steroids are used appropriately for a relatively short time they are very, very safe
- After all, they are basically a natural substance already found in the body
- Be aware of body weight when dosing

## WHO DOESN'T GET ORALS, OR GETS THEM VERY, VERY CAREFULLY

- Diabetics
- Patients with stomach problems / ulcers
- Patients with active infection
- Pregnant women



## WHAT CAN THEY DO THAT'S BAD?

- Almost nothing in the short term!
- Increase Na<sup>+</sup>, decreased K leading to fluid retention
- Hypertension
- Elevate blood glucose levels
- Stomach pain and ulcers
- Insomnia, euphoria, psychosis
- Thin skin / bruising
- Osteoporosis
- Increased ICP
- PSC's (more than topicals)
- Increased IOP (less than topicals)



## WHAT CAN THEY INTERACT WITH?

- Screw up glucose control
- ASA, Coumadin
- Digoxin
- Some antibiotics, anti-seizure meds, anti-TB meds (TB itself is a strong relative contraindication)



## WHAT DO THEY DO THAT'S GOOD?

- Duh!.....they decrease inflammation and therefore inflammatory sequelae



## WHAT CAN WE USE ORAL STEROIDS FOR IN EYE CARE?

- Contact dermatitis / allergic response of the eye lids
- Reaction to insect bite or sting on the eye lids
- Recalcitrant CME
- Recalcitrant uveitis, especially bilateral or vitritis
- Choroiditis / retinitis
- Scleritis

## USES OF ORALS IN EYE CARE

- Myasthenia Gravis
- Inflammatory orbital pseudotumor
- Thyroid eye disease / Grave's ophthalmopathy
- Optic neuritis (but not by themselves!)
- GCA
- DLK post LASIK (in conjunction with topicals)

## OCULAR SIDE EFFECTS OF ORAL STEROIDS

- These are well known.....PSC's and increased IOP
- IOP increases are rare, but can occur with very long term use
- PSC's are not rare!
- 10 mg per day or less for one year or less has almost no chance of PSC formation
- 16 mg per day or more for several years has a 75% chance of PSC formation
- Overall, general population has a .5% chance of PSC development while those on long term oral steroids have a 30% prevalence (across doses)

## ORAL STEROIDS



- Oral steroids are generally prescribed in one of two ways.....
- 1) Medrol dose pack (methylprednisolone)
- 2) Prednisone 10mg tablets

## COMPARISONS

- When it comes to suppressing the HPA (hypothalamic-pituitary-adrenal) axis.....
- 25mg Cortisone = 20mg Hydrocortisone = 5mg Prednisone = 4mg Triamcinolone = 4mg Methylprednisone = .75mg Dexamethasone
- Potency essentially follows this order but in reverse
- Body produces an amount of cortisone that equals 5mg of prednisone per day



## MEDROL DOSE PACK

- Available in different strengths
- Most commonly used is a package of 21, 4 mg tablets( 2 mg is available)
- Six are taken the first day, then one less each day thereafter (6-5-4-3-2-1 = 21 tablets)
- Self tapering and little to no suppression of the HPA axis
- In eye care, really only strong enough and long lasting enough for treatment of lid reactions

## PREDNISON\*

- Most common dosing is to give the desired amount in 10 mg tablets (need 40 mg, take 4 pills)
- Is available in 1, 2.5, 5, 10, 20, and 50 mg tablets
- Best choice for most of our desired uses in eye care
- Potent and flexible

## DOSING

- Up to 60 mg, take entire dose in the morning
- Over this amount take  $\frac{1}{2}$  in morning,  $\frac{1}{2}$  in evening
- As previously mentioned, Medrol dose pack self tapers
- With prednisone, after relatively short course at full desired strength (say 2-3 weeks or so), taper by ten milligrams every other day

## DOSING

- An alternative approach is to give twice the desired dose every other day then don't taper. Only for short term use
- Theory is that anti-inflammatory properties remain high but suppression of HPA axis is much, much less
- For long term use taper must be very slow
- As OD's we rarely would be involved in the long term prescription of oral steroids

THE END!



- Questions or comments: [brsutton@indiana.edu](mailto:brsutton@indiana.edu)